


## Sacramento River Hydrologic Region



**Figure 33 Sacramento River Hydrologic Region**

## Basins and Subbasins of the Sacramento River Hydrologic Region

Basin/subbasins	Basin name	Basin/subbasins	Basin name
5-1	Goose Lake Valley	5-30	Lower Lake Valley
5-1.01	Lower Goose Lake Valley	5-31	Long Valley
5-1.02	Fandango Valley	5-35	Mccloud Area
5-2	Alturas Area	5-36	Round Valley
5-2.01	South Fork Pitt River	5-37	Toad Well Area
5-2.02	Warm Springs Valley	5-38	Pondosa Town Area
5-3	Jess Valley	5-40	Hot Springs Valley
5-4	Big Valley	5-41	Egg Lake Valley
5-5	Fall River Valley	5-43	Rock Prairie Valley
5-6	Redding Area	5-44	Long Valley
5-6.01	Bowman	5-45	Cayton Valley
5-6.02	Rosewood	5-46	Lake Britton Area
5-6.03	Anderson	5-47	Goose Valley
5-6.04	Enterprise	5-48	Burney Creek Valley
5-6.05	Millville	5-49	Dry Burney Creek Valley
5-6.06	South Battle Creek	5-50	North Fork Battle Creek
5-7	Lake Almanor Valley	5-51	Butte Creek Valley
5-8	Mountain Meadows Valley	5-52	Gray Valley
5-9	Indian Valley	5-53	Dixie Valley
5-10	American Valley	5-54	Ash Valley
5-11	Mohawk Valley	5-56	Yellow Creek Valley
5-12	Sierra Valley	5-57	Last Chance Creek Valley
5-12.01	Sierra Valley	5-58	Clover Valley
5-12.02	Chilcoot	5-59	Grizzly Valley
5-13	Upper Lake Valley	5-60	Humbug Valley
5-14	Scotts Valley	5-61	Chrome Town Area
5-15	Big Valley	5-62	Elk Creek Area
5-16	High Valley	5-63	Stonyford Town Area
5-17	Burns Valley	5-64	Bear Valley
5-18	Coyote Valley	5-65	Little Indian Valley
5-19	Collayomi Valley	5-66	Clear Lake Cache Formation
5-20	Berryessa Valley	5-68	Pope Valley
5-21	Sacramento Valley	5-86	Joseph Creek
5-21.50	Red Bluff	5-87	Middle Fork Feather River
5-21.51	Corning	5-88	Stony Gorge Reservoir
5-21.52	Colusa	5-89	Squaw Flat
5-21.53	Bend	5-90	Funks Creek
5-21.54	Antelope	5-91	Antelope Creek
5-21.55	Dye Creek	5-92	Blanchard Valley
5-21.56	Los Molinos	5-93	North Fork Cache Creek
5-21.57	Vina	5-94	Middle Creek
5-21.58	West Butte	5-95	Meadow Valley
5-21.59	East Butte		
5-21.60	North Yuba		
5-21.61	South Yuba		
5-21.62	Sutter		
5-21.64	North American		
5-21.65	South American		
5-21.66	Solano		
5-21.67	Yolo		
5-21.68	Capay Valley		

## Description of the Region

The Sacramento River HR covers approximately 17.4 million acres (27,200 square miles). The region includes all or large portions of Modoc, Siskiyou, Lassen, Shasta, Tehama, Glenn, Plumas, Butte, Colusa, Sutter, Yuba, Sierra, Nevada, Placer, Sacramento, El Dorado, Yolo, Solano, Lake, and Napa counties (Figure 33). Small areas of Alpine and Amador counties are also within the region. Geographically, the region extends south from the Modoc Plateau and Cascade Range at the Oregon border, to the Sacramento-San Joaquin Delta. The Sacramento Valley, which forms the core of the region, is bounded to the east by the crest of the Sierra Nevada and southern Cascades and to the west by the crest of the Coast Range and Klamath Mountains. Other significant features include Mount Shasta and Lassen Peak in the southern Cascades, Sutter Buttes in the south central portion of the valley, and the Sacramento River, which is the longest river system in the State of California with major tributaries the Pit, Feather, Yuba, Bear and American rivers. The region corresponds approximately to the northern half of RWQCB 5. The Sacramento metropolitan area and surrounding communities form the major population center of the region. With the exception of Redding, cities and towns to the north, while steadily increasing in size, are more rural than urban in nature, being based in major agricultural areas. The 1995 population of the entire region was 2.372 million.

The climate in the northern, high desert plateau area of the region is characterized by cold snowy winters with only moderate precipitation and hot dry summers. This area depends on adequate snowpack to provide runoff for summer supply. Annual precipitation ranges from 10 to 20 inches. Other mountainous areas in the northern and eastern portions of the region have cold wet winters with large amounts of snow, which typically provide abundant runoff for summer supplies. Annual precipitation ranges from 40 to more than 80 inches. Summers are generally mild in these areas. The Coast Range and southern Klamath Mountains receive copious amounts of precipitation, but most of the runoff flows to the coast in the North Coastal drainage. Sacramento Valley comprises the remainder of the region. At a much lower elevation than the rest of the region, the valley has mild winters with moderate precipitation. Annual precipitation varies from about 35 inches in Redding to about 18 inches in Sacramento. Summers in the valley are hot and dry.

Most of the mountainous portions of the region are heavily forested and sparsely populated. Three major national forests (Mendocino, Trinity, and Shasta) make up the majority of lands in the Coast Range, southern Klamath Mountains, and the southern Cascades; these forests and the region's rivers and lakes provide abundant recreational opportunities. In the few mountain valleys with arable land, alfalfa, grain and pasture are the predominant crops. In the foothill areas of the region, particularly adjacent to urban centers, suburban to rural housing development is occurring along major highway corridors. This development is leading to urban sprawl and is replacing the former agricultural production on those lands. In the Sacramento Valley, agriculture is the largest industry. Truck, field, orchard, and rice crops are grown on approximately 2.1 million acres. Rice represents about 23 percent of the total irrigated acreage.

The Sacramento River HR is the main water supply for much of California's urban and agricultural areas. Annual runoff in the HR averages about 22.4 maf, which is nearly one-third of the State's total natural runoff. Major water supplies in the region are provided through surface storage reservoirs. The two largest surface water projects in the region are USBR's Shasta Lake (Central Valley Project) on the upper Sacramento River and Lake Oroville (DWR's State Water Project) on the Feather River. In all, there are more than 40 major surface water reservoirs in the region. Municipal, industrial, and agricultural supplies to the region are about 8 maf, with groundwater providing about 2.5 maf of that total. Much of the remainder of the runoff goes to dedicated natural flows, which support various environmental requirements, including in-stream fishery flows and flushing flows in the Delta.

## Groundwater Development

Groundwater provides about 31 percent of the water supply for urban and agricultural uses in the region, and has been developed in both the alluvial basins and the hard rock uplands and mountains. There are 88 basins/subbasins delineated in the region. These basins underlie 5.053 million acres (7,900 square miles), about 29 percent of the entire region. The reliability of the groundwater supply varies greatly. The Sacramento Valley is recognized as one of the foremost groundwater basins in the State, and wells developed in the sediments of the valley provide excellent supply to irrigation, municipal, and domestic uses. Many of the mountain valleys of the region also provide significant groundwater supplies to multiple uses.

Geologically, the Sacramento Valley is a large trough filled with sediments having variable permeabilities; as a result, wells developed in areas with coarser aquifer materials will produce larger amounts of water than wells developed in fine aquifer materials. In general, well yields are good and range from one-hundred to several thousand gallons per minute. Because surface water supplies have been so abundant in the valley, groundwater development for agriculture primarily supplement the surface supply. With the changing environmental laws and requirements, this balance is shifting to a greater reliance on groundwater, and conjunctive use of both supplies is occurring to a greater extent throughout the valley, particularly in drought years. Groundwater provides all or a portion of municipal supply in many valley towns and cities. Redding, Anderson, Chico, Marysville, Sacramento, Olivehurst, Wheatland, Willows, and Williams rely to differing degrees on groundwater. Red Bluff, Corning, Woodland, Davis, and Dixon are completely dependent on groundwater. Domestic use of groundwater varies, but in general, rural unincorporated areas rely completely on groundwater.

In the mountain valleys and basins with arable land, groundwater has been developed to supplement surface water supplies. Most of the rivers and streams of the area have adjudicated water rights that go back to the early 1900s, and diversion of surface water has historically supported agriculture. Droughts and increased competition for supply have led to significant development of groundwater for irrigation. In some basins, the fractured volcanic rock underlying the alluvial fill is the major aquifer for the area. In the rural mountain areas of the region, domestic supplies come almost entirely from groundwater. Although a few mountain communities are supplied in part by surface water, most rely on groundwater. These groundwater supplies are generally quite reliable in areas that have sufficient aquifer storage or where surface water replenishes supply throughout the year. In areas that depend on sustained runoff, water levels can be significantly depleted in drought years and many old, shallow wells can be dewatered. During 2001, an extreme drought year on the Modoc Plateau, many well owners experienced problems with water supply.

Groundwater development in the fractured rocks of the foothills of the southern Cascades and Sierra Nevada is fraught with uncertainty. Groundwater supplies from fractured rock sources are highly variable in terms of water quantity and water quality and are an uncertain source for large-scale residential development. Originally, foothill development relied on water supply from springs and river diversions with flumes and ditches for conveyance that date back to gold mining era operations. Current development is primarily based on individual private wells, and as pressures for larger scale development increase, questions about the reliability of supply need to be addressed. Many existing foothill communities have considerable experience with dry or drought year shortages. In Butte County residents in Cohasset, Forest Ranch, and Magalia have had to rely on water brought up the ridges in tanker trucks. The suggested answer has been the development of regional water supply projects. Unfortunately, the area's development pattern of small, geographically dispersed population centers does not lend itself to the kind of financial base necessary to support such projects.

## Groundwater Quality

Groundwater quality in the Sacramento River HR is generally excellent. However, there are areas with local groundwater problems. Natural water quality impairments occur at the north end of the Sacramento Valley in the Redding subbasin, and along the margins of the valley and around the Sutter Buttes, where Cretaceous-age marine sedimentary rocks containing brackish to saline water are near the surface. Water from the older underlying sediments mixes with the fresh water in the younger alluvial aquifer and degrades the quality. Wells constructed in these areas typically have high TDS. Other local natural impairments are moderate levels of hydrogen sulfide in groundwater in the volcanic and geothermal areas in the western portion of the region. In the Sierra foothills, there is potential for encountering uranium and radon-bearing rock or sulfide mineral deposits containing heavy metals. Human-induced impairments are generally associated with individual septic system development in shallow unconfined portions of aquifers or in fractured hard rock areas where insufficient soil depths are available to properly leach effluent before it reaches the local groundwater supply.

### Water Quality in Public Supply Wells

From 1994 through 2000, 1,356 public supply water wells were sampled in 51 of the 88 basins and subbasins in the Sacramento River HR. Samples analyzed indicate that 1,282 wells, or 95 percent, met the state primary MCLs for drinking water. Seventy-four wells, or 5 percent, have constituents that exceed one or more MCL. Figure 34 shows the percentages of each contaminant group that exceeded MCLs in the 74 wells.




Figure 34 MCL exceedances in public supply wells in the Sacramento River Hydrologic Region

Table 25 lists the three most frequently occurring contaminants in each of the six contaminant groups and shows the number of wells in the HR that exceeded the MCL for those contaminants.

**Table 25 Most frequently occurring contaminants by contaminant group in the Sacramento River Hydrologic Region**

Contaminant group	Contaminant - # of wells	Contaminant - # of wells	Contaminant - # of wells
Inorganics – Primary	Cadmium – 4	Chromium (Total) – 3	3 tied at 2
Inorganics – Secondary	Manganese – 221	Iron – 166	Specific Conductance – 3
Radiological	Gross Alpha – 4		
Nitrates	Nitrate (as NO <sub>3</sub> ) – 22	Nitrate + Nitrite – 5	Nitrate Nitrogen (NO <sub>3</sub> -N) – 2
Pesticides	Di(2-Ethylhexyl)phthalate – 4		
VOCs/SVOCs	PCE – 11	TCE – 7	Benzene – 4

PCE = Tetrachloroethylene

TCE = Trichloroethylene

VOC = Volatile Organic Compounds

SVOC = Semivolatile Organic Compound

### Changes from Bulletin 118-80

Some modifications from the groundwater basins presented in Bulletin 118-80 are incorporated in this report. These are listed in Table 26.

**Table 26 Modifications since Bulletin 118-80 of groundwater basins and subbasins in Sacramento River Hydrologic Region**

Basin name	New number	Old number
Fandango Valley	5-1.02	5-39
Bucher Swamp Valley	deleted	5-42
Modoc Plateau Recent Volcanic Areas	deleted	5-32
Modoc Plateau Pleistocene Volcanic Areas	deleted	5-33
Mount Shasta Area	deleted	5-34
Sacramento Valley Eastside Tuscan Formation Highlands	deleted	5-55
Clear Lake Pleistocene Volcanics	deleted	5-67

No additional basins were assigned to the Sacramento River HR in this revision. However, four basins have been divided into subbasins. Goose Lake Valley Groundwater Basin (5-1) has been subdivided into two subbasins, Fandango Valley (5-39) was modified to be a subbasin of Goose Lake Valley. Redding Area Groundwater Basin has been subdivided into six subbasins, Sierra Valley Groundwater Basin has been subdivided into two subbasins, and the Sacramento Valley Groundwater Basin has been subdivided into 18 subbasins.

There are several deletions of groundwater basins from Bulletin 118-80. Bucher Swamp Valley Basin (5-42) was deleted due to a thin veneer of alluvium over rock. Modoc Plateau Recent Volcanic Areas (5-32), Modoc Plateau Pleistocene Volcanic Areas (5-33), Mount Shasta Area (5-34), Sacramento Valley Eastside Tuscan Formation Highlands (5-55), and Clear Lake Pleistocene Volcanics (5-67) are volcanic aquifers and were not assigned basin numbers in this bulletin. These are considered to be groundwater source areas as discussed in Chapter 6.



Table 27 Sacramento River Hydrologic Region groundwater data

Basin/Subbasin	Basin Name	Area (acres)	Groundwater Budget Type	Well Yields (gpm)		Types of Monitoring			TDS (mg/L)	
				Maximum	Average	Levels	Quality	Title 22	Average	Range
5-1	GOOSE LAKE VALLEY									
5-1.01	LOWER GOOSE LAKE	36,000	B	-	400	9	9	-	183	68 - 528
5-1.02	FANDANGO VALLEY	18,500	B	2,000	-	3	-	-	-	-
5-2	ALTURAS AREA									
5-2.01	SOUTH FORK PITT RIVER	114,000	B	5,000	1,075	9	-	8	-	-
5-2.02	WARM SPRINGS VALLEY	68,000	B	400	314	3	-	11	-	-
5-3	JESS VALLEY	6,700	B	-	3,000	-	-	-	-	-
5-4	BIG VALLEY	92,000	B	4,000	880	19	9	10	260	141 - 633
5-5	FALL RIVER VALLEY	54,800	B	1,500	266	16	7	3	174	115 - 232
5-6	REDDING AREA									
5-6.01	BOWMAN	85,330	B	2,000	589	8	2	13	-	70 - 247
5-6.02	ROSEWOOD	45,320	B	-	-	4	-	-	-	118 - 218
5-6.03	ANDERSON	98,500	B	1,800	46	11	10	69	194	109-320
5-6.04	ENTERPRISE	60,900	B	700	266	11	3	43	-	160 - 210
5-6.05	MILLVILLE	67,900	B	500	254	6	5	4	140	-
5-6.06	SOUTH BATTLE CREEK	32,300	B	-	-	0	0	0	360	-
5-7	LAKE ALMANOR VALLEY	7,150	B	-	-	10	4	4	105	53 - 260
5-8	MOUNTAIN MEADOWS VALLEY	8,150	B	-	-	-	-	-	-	-
5-9	INDIAN VALLEY	29,400	B	-	-	-	4	9	-	-
5-10	AMERICAN VALLEY	6,800	B	40	40	-	4	11	-	-
5-11	MOHAWK VALLEY	19,000	B	-	500	1	2	15	248	210 - 285
5-12	SIERRA VALLEY									
5-12.01	SIERRA VALLEY	117,700	B	1,500	640	34	15	9	312	110 - 1,620
5-12.02	CHILCOOT	7,550	B	-	-	15	-	8	-	-
5-13	UPPER LAKE VALLEY	7,260	B	900	302	12	3	6	-	-
5-14	SCOTTS VALLEY	7,320	B	1,200	171	9	1	9	158	140 - 175
5-15	BIG VALLEY	24,210	B	1,470	475	49	11	7	535	270 - 790
5-16	HIGH VALLEY	2,360	B	100	37	5	2	-	598	480 - 745
5-17	BURNS VALLEY	2,900	B	-	30	1	5	-	335	280 - 455
5-18	COYOTE VALLEY	6,530	B	800	446	6	3	3	288	175 - 390
5-19	COLLAYOMI VALLEY	6,500	B	1,000	121	10	4	3	202	150 - 255
5-20	BERRYESSA VALLEY	1,400	C	-	-	0	-	0	-	-
5-21	SACRAMENTO VALLEY									
5-21.50	RED BLUFF	266,750	B	1,200	363	30	10	56	207	120 - 500
5-21.51	CORNING	205,640	B	3,500	977	29	7	30	286	130 - 490
5-21.52	COLUSA	918,380	B	5,600	984	98	30	134	391	120 - 1,220
5-21.53	BEND	20,770	B	-	275	0	3	9	-	334-360
5-21.54	ANTELOPE	18,710	B	800	575	4	5	22	296	-
5-21.55	DYE CREEK	27,730	B	3,300	890	8	1	3	240	159 - 396
5-21.56	LOS MOLINOS	33,170	B	1,000	500	3	3	9	217	-
5-21.57	VINA	125,640	B	3,850	1,212	23	5	69	285	48 - 543
5-21.58	WEST BUTTE	181,600	B	4,000	1,833	32	8	36	293	130 - 676

Table 27 Sacramento River Hydrologic Region groundwater data (continued)

Basin/Subbasin	Basin Name	Area (acres)	Groundwater Budget Type	Well Yields (gpm)		Types of Monitoring			TDS (mg/L)	
				Maximum	Average	Levels	Quality	Title 22	Average	Range
5-21.59	EAST BUTTE	265,390	B	4,500	1,019	43	4	44	235	122 - 570
5-21.60	NORTH YUBA	100,400	C	4,000	-	21	-	32	-	-
5-21.61	SOUTH YUBA	107,000	C	4,000	1,650	56	-	6	-	-
5-21.62	SUTTER	234,000	C	-	-	34	-	115	-	-
5-21.64	NORTH AMERICAN	351,000	A	-	800	121	-	339	300	150 - 1,000
5-21.65	SOUTH AMERICAN	248,000	C	-	-	105	-	247	221	24 - 581
5-21.66	SOLANO	425,000	C	-	-	123	23	136	427	150 - 880
5-21.67	YOLO	226,000	B	4,000+	1,000	127	20	185	880	480 - 2,060
5-21.68	CAPAY VALLEY	25,000	C	-	-	11	-	3	-	-
5-30	LOWER LAKE VALLEY	2,400	B	100	37	-	3	5	568	290 - 1,230
5-31	LONG VALLEY	2,600	B	100	63	-	-	-	-	-
5-35	MCCLOUD AREA	21,320	B	-	380	-	-	1	-	-
5-36	ROUND VALLEY	7,270	B	2,000	800	2	-	-	-	148 - 633
5-37	TOAD WELL AREA	3,360	B	-	-	-	-	-	-	-
5-38	PONDOSA TOWN AREA	2,080	B	-	-	-	-	-	-	-
5-40	HOT SPRINGS VALLEY	2,400	B	-	-	-	-	-	-	-
5-41	EGG LAKE VALLEY	4,100	B	-	20	-	-	-	-	-
5-43	ROCK PRAIRIE VALLEY	5,740	B	-	-	-	-	-	-	-
5-44	LONG VALLEY	1,090	B	-	-	-	-	-	-	-
5-45	CAYTON VALLEY	1,300	B	-	400	-	-	-	-	-
5-46	LAKE BRITTON AREA	14,060	B	-	-	-	-	2	-	-
5-47	GOOSE VALLEY	4,210	B	-	-	-	-	-	-	-
5-48	BURNEY CREEK VALLEY	2,350	B	-	-	-	-	2	-	-
5-49	DRY BURNEY CREEK VALLEY	3,070	B	-	-	-	-	-	-	-
5-50	NORTH FORK BATTLE CREEK VALLEY	12,760	B	-	-	-	-	3	-	-
5-51	BUTTE CREEK VALLEY	3,230	B	-	-	-	-	-	-	-
5-52	GRAYS VALLEY	5,440	B	-	-	-	-	-	-	-
5-53	DIXIE VALLEY	4,870	B	-	-	-	-	-	-	-
5-54	ASH VALLEY	6,010	B	3,000	2,200	-	-	-	-	-
5-56	YELLOW CREEK VALLEY	2,310	B	-	-	-	-	-	-	-
5-57	LAST CHANCE CREEK VALLEY	4,660	B	-	-	-	-	-	-	-
5-58	CLOVER VALLEY	16,780	B	-	-	-	-	-	-	-
5-59	GRIZZLY VALLEY	13,400	B	-	-	-	-	1	-	-
5-60	HUMBUG VALLEY	9,980	B	-	-	-	-	8	-	-
5-61	CHROME TOWN AREA	1,410	B	-	-	-	-	-	-	-
5-62	ELK CREEK AREA	1,440	B	-	-	-	-	-	-	-
5-63	STONYFORD TOWN AREA	6,440	B	-	-	-	-	-	-	-
5-64	BEAR VALLEY	9,100	B	-	-	-	-	-	-	-
5-65	LITTLE INDIAN VALLEY	1,270	B	-	-	-	-	-	-	-
5-66	CLEAR LAKE CACHE FORMATION	30,000	B	-	52	-	-	4	-	-
5-68	POPE VALLEY	7,180	C	-	-	-	-	1	-	-
5-86	JOSEPH CREEK	4,450	B	-	-	-	-	-	-	-

Table 27 Sacramento River Hydrologic Region groundwater data (continued)

Basin/Subbasin	Basin Name	Area (acres)	Groundwater Budget Type	Well Yields (gpm)		Types of Monitoring			TDS (mg/L)	
				Maximum	Average	Levels	Quality	Title 22	Average	Range
5-87	MIDDLE FORK FEATHER RIVER	4,340	B	-	-	-	-	2	-	-
5-88	STONY GORGE RESERVOIR	1,070	B	-	-	-	-	-	-	-
5-89	SQUAW FLAT	1,300	C	-	-	-	-	-	-	-
5-90	FUNKS CREEK	3,000	C	-	-	-	-	-	-	-
5-91	ANTELOPE CREEK	2,040	B	-	-	-	-	-	-	-
5-92	BLANCHARD VALLEY	2,200	B	-	-	-	-	-	-	-
5-93	NORTH FORK CACHE CREEK	3,470	C	-	-	-	-	-	-	-
5-94	MIDDLE CREEK	700	B	-	75	-	-	1	-	-
5-95	MEADOW VALLEY	5,730	B	-	-	-	-	1	-	-

gpm - gallons per minute

mg/L - milligram per liter

TDS -total dissolved solids

## Sacramento Valley Groundwater Basin, East Butte Subbasin

- Groundwater Basin Number: 5-21.59
- County: Butte, Sutter
- Surface Area: 265,390 acres (415 square miles)

### Basin Boundaries and Hydrology

The East Butte Subbasin is the portion of the Sacramento Valley Groundwater Basin bounded on the west and northwest by Butte Creek, on the northeast by the Cascade Ranges, on the southeast by the Feather River and the south by the Sutter Buttes. The northeast boundary along the Cascade Ranges is primarily a geographic boundary with some groundwater recharge occurring beyond that boundary. The subbasin is contiguous with the West Butte Subbasin at depth. Annual precipitation is approximately 18 inches in the valley increasing to 27 inches towards the eastern foothills.

### Hydrogeologic Information

#### *Water-Bearing Formations*

The East Butte aquifer system is comprised of deposits of late Tertiary to Quaternary age. The Quaternary deposits include Holocene stream channel deposits and basin deposits, Pleistocene deposits of the Modesto and Riverbank formations, and Sutter Buttes alluvium. The Tertiary deposits include the Tuscan and Laguna formations.

**Holocene Stream Channel Deposits.** These deposits consist of unconsolidated gravel, sand, silt and clay derived from the erosion, reworking, and deposition of adjacent Quaternary stream terrace alluvial deposits. The thickness varies from 1- to 80-feet (Helley and Harwood 1985). These deposits represent the upper part of the unconfined zone of the aquifer and are moderately-to-highly permeable; however, the thickness and areal extent of the deposits limit the water-bearing capability.

**Holocene Basin Deposits.** These deposits are the result of sediment-laden floodwaters that rose above the natural levees of streams and rivers to spread across low-lying areas. They consist primarily of silts and clays and may be locally interbedded with stream channel deposits. These deposits result from deposition from erosion from portions of the Cascade Ranges to the Sutter Buttes. Thickness of the deposits range to 150 feet (DWR 2000). These deposits have low permeability and generally yield low quantities of water to wells. The quality of groundwater produced from the basin deposits is often poor (USBR 1960).

**Pleistocene Modesto Formation.** The Modesto Formation in this subbasin consists of poorly indurated gravel and cobbles with sand, silt, and clay derived from reworking and deposition of the Tuscan Formation, Laguna Formation, and the Riverbank Formation. Surface exposure of the formation is west of the Feather River extending from south of the Thermalito Afterbay to the southern subbasin boundary. The formation may extend across the entire subbasin, underlying basin deposits, with thicknesses ranging from 50- to 150-feet (DWR 2000).

**Pleistocene Riverbank Formation.** These older terrace deposits consist of poorly-to-highly permeable pebble and small cobble gravels interlensed with reddish clay sands and silt. Surface exposure of the Riverbank Formation is primarily south and west of the Thermalito Afterbay. The formation may extend across the entire subbasin, underlying basin and Modesto deposits, with thicknesses ranging from 50- to 200-feet (Helley and Harwood 1985).

**Pleistocene Sutter Butte Alluvium.** In the southern portion of the subbasin, alluvium of the Sutter Buttes is observed in the subsurface and may range in thickness up to 600 feet (DWR 2000). The fan deposits forming the apron around the buttes consist largely of gravel, sand, silt and clay and may extend up to 15 miles north of the Sutter Buttes and westerly beyond the Sacramento River. Utility pump test records show the average well yield for that formation to be approximately 2300 gallons per minute with an average specific capacity of 64.

**Pliocene Tuscan Formation.** The Tuscan Formation is composed of a series of volcanic mudflows, tuff breccia, tuffaceous sandstone and volcanic ash layers. Thickness of the formation is estimated to be 800 feet (DWR 2000). The formation is described as four separate but lithologically similar units, A through D (with Unit A being the oldest), which in some areas are separated by layers of thin tuff or ash units (Helley and Harwood 1985). Units A, B, and C are found within the subsurface in the northern part of the subbasin and Units A and B are found in the southern part of the subbasin. Surface exposures of Units B and C are located in the foothills at the far eastern extents of the subbasin.

Unit A is the oldest water bearing unit of the formation and is characterized by the presence of metamorphic clasts within interbedded lahars, volcanic conglomerate, volcanic sandstone and siltstone.

Unit B is composed of fairly equal distribution of lahars, tuffaceous sandstone, and conglomerate. Unit C consists of massive mudflow or lahar deposits with some interbedded volcanic conglomerate and sandstone. In the subsurface, these low permeability lahars form thick, confining layers for groundwater contained in the more permeable sediments if Unit B.

**Pliocene Laguna Formation.** The Laguna Formation consists of interbedded alluvial sand, gravel, and silt deposits which are moderately consolidated and poorly-to-well cemented. The Laguna is compacted and generally has a low-to-moderate permeability, except in scattered gravels in the upper portion. The formation yields moderate quantities of water to wells along the eastern margin of the valley. Wells of higher capacity generally tap underlying Tuscan deposits.

Surface exposures of the Laguna appear along the eastern margin of the subbasin in the vicinity of the Thermalito Afterbay and extend westerly in the subsurface. The lateral extent of the formation is unknown. The thickness of the formation is difficult to determine because the base of the unit is rarely exposed. Estimates of maximum thickness range from 180 feet (Helley and Harwood 1985) to 1,000 feet (Olmsted and Davis 1961). Geologic cross sections developed by California Department of Water

Resources estimate the thickness to be approximately 500 feet (DWR 2000). Wells completed in the formation yield only moderate quantities of water.

### ***Groundwater Level Trends***

As part of a groundwater inventory analysis prepared for Butte County, the portion of the East Butte Subbasin located within Butte County was evaluated for seasonal and long-term changes in groundwater levels for confined and composite portions of the aquifer systems (DWR 2001).

For wells constructed in confined and composite portions of the aquifer, the increased use of groundwater in the northern portion of the subbasin has resulted in wide seasonal fluctuations in groundwater levels. In the northern portion of the subbasin, composite well fluctuations (composite wells are monitoring wells that represent groundwater levels that combine confined and unconfined portions of the aquifer system) average about 15 feet during normal years and 30- to 40- feet during drought years. Annual groundwater fluctuations in the confined and semi-confined aquifer system ranges from 15- to 30- feet during normal years.

In the portion of the subbasin located within the southern part of Butte County, groundwater level fluctuations for composite wells average about 4 feet during normal years and up to 10 feet during drought years. The groundwater fluctuations for wells constructed in the confined and semi-confined aquifer system average 4 feet during normal years and up to 5 feet during drought years.

### ***Recharge Areas***

Localized fluctuations in groundwater levels are observed just south of the Thermalito Afterbay due to the recharging of groundwater from this surface water system (DWR 2001).

### ***Groundwater Storage***

The storage capacity of the subbasin was estimated based on estimates of specific yield for the Sacramento Valley as developed in DWR (1978). Estimates of specific yield, determined on a regional basis, were used to obtain a weighted specific yield conforming to the subbasin boundary. The estimated specific yield for the East Butte Subbasin is 5.9 percent. The estimated storage capacity to a depth of 200 feet is approximately 3,128,959 acre-feet.

### ***Groundwater Budget (Type B)***

Estimates of groundwater extraction are based on surveys conducted by the California Department of Water Resources during 1993 and 1997. Surveys included land use and sources of water. Estimates of groundwater extraction for agricultural; municipal and industrial; and environmental wetland uses are 104,000, 75,500 and 1,300 acre-feet respectively. Deep percolation of applied water is estimated to be 126,000 acre-feet.

### ***Groundwater Quality***

**Characterization.** Calcium-magnesium bicarbonate and magnesium-calcium bicarbonate waters are the predominant groundwater water types in the

subbasin. Magnesium bicarbonate waters occur locally near Biggs-Gridley, south and east to the Feather River. Total dissolved solids range from 122- to 570-mg/L, averaging 235 mg/L (DWR unpublished data).

**Impairments.** Localized high concentrations of manganese, iron, magnesium, total dissolved solids, conductivity, ASAR, and calcium occur within the subbasin.

### Water Quality in Public Supply Wells

Constituent Group <sup>1</sup>	Number of wells sampled <sup>2</sup>	Number of wells with a concentration above an MCL <sup>3</sup>
Inorganics – Primary	30	1
Radiological	25	0
Nitrates	32	2
Pesticides	16	0
VOCs and SVOCs	19	0
Inorganics – Secondary	30	3

<sup>1</sup> A description of each member in the constituent groups and a generalized discussion of the relevance of these groups are included in *California's Groundwater – Bulletin 118* by DWR (2003).

<sup>2</sup> Represents distinct number of wells sampled as required under DHS Title 22 program from 1994 through 2000.

<sup>3</sup> Each well reported with a concentration above an MCL was confirmed with a second detection above an MCL. This information is intended as an indicator of the types of activities that cause contamination in a given basin. It represents the water quality at the sample location. It does not indicate the water quality delivered to the consumer. More detailed drinking water quality information can be obtained from the local water purveyor and its annual Consumer Confidence Report.

### Well Production characteristics

Well yields (gal/min)		
Irrigation	Range: 0 – 4500	Average: 1839 (37 Well Completion Reports)
Utility pump test records for the East Butte Subbasin show well yields ranging from a low of 65 gpm to a high of 5,459 gpm with an average yield of 1,602 gpm (DWR 2001).		
Total depths (ft)		
Domestic	Range: 25 – 639	Average: 101 (1477 Well Completion Reports)
Irrigation	Range: 35 – 983	Average: 285 (699 Well Completion Reports)

### Active Monitoring Data

Agency	Parameter	Number of wells /measurement frequency
DWR	Groundwater levels	43 wells semi-annually
DWR	Miscellaneous water quality	4 wells biennially
Department of Health Services	Miscellaneous water quality	44

## Basin Management

---

Groundwater management: Butte County adopted a groundwater management ordinance in 1996.

### Water agencies

Public	Butte Basin Water Users Association, Biggs-West Gridley WD, Butte WD, Durham ID, City of Biggs, City of Gridley, Oroville-Wyandotte ID, Richvale ID, Thermalito ID, and Western Canal WD.
Private	North Burbank Public Utility District.

---

## Selected References

- California Department of Water Resources. 1978. Evaluation of Groundwater Resources: Sacramento Valley. Department of Water Resources in cooperation with the United States Geological Survey. Appendix A. Bulletin 118-6.
- California Department of Water Resources. 2001. Butte County Groundwater Inventory Analysis. Draft Report. Northern District.
- California Department of Water Resources. 2000. Geology and Hydrogeology of the Freshwater Bearing Aquifer Systems of the Northern Sacramento Valley, California. In Progress.
- Helley EJ, Harwood DS. 1985. Geologic Map of the Late Cenozoic Deposits of the Sacramento Valley and Northern Sierran Foothills, California. Map MF-1790.
- Olmsted FH, Davis GH. 1961. Geologic Features and Ground Water Storage Capacity of the Sacramento Valley, California. USGS. Water Supply Paper 1497.
- United States Bureau of Reclamation (USBR), 1960. Tehama-Colusa Service Area Geology and Groundwater Resources Appendix.

## Bibliography

- Bailey EH. 1966. Geology of Northern California. California Division of Mines and Geology. Bulletin 190.
- Berkstressor CF. 1973. Base of Fresh Water in the Sacramento Valley and Sacramento-San Joaquin Delta, California. U.S. Geological Survey in Cooperation with California Department of Water Resources.
- Bertoldi GT, Johnson RH, Evenson KD. 1991. Groundwater in the Central Valley, California - A Summary Report. Regional Aquifer System Analysis--Central Valley, California. USGS. Professional Paper 1401-A.
- Beyer LA. 1993. Sacramento Basin Province. USGS.
- Bryan K. 1923. Geology and Ground-water Resources of Sacramento Valley, California. USGS. 495.
- Busacca AJ, Singer MJ, Verosub KL. 1989. Late Cenozoic Stratigraphy of the Feather and Yuba Rivers Area, California, with a Section on Soil Development in Mixed Alluvium at Honcut Creek. USGS.
- California Department of Pesticide Regulation. 1993. Sampling for Pesticide Residues in California Well Water, 1993 Well Inventory Database. California Environmental Protection Agency.
- California Department of Water Resources. 1958. Ground Water Conditions in Central and Northern California 1957-58. California Department of Water Resources. Bulletin 77-58.



- California Department of Water Resources. 1960. Northeastern Counties Investigation. California Department of Water Resources. Bulletin 58.
- California Department of Water Resources. 1964. Groundwater Conditions in Central and Northern California, 1961-62. California Department of Water Resources.
- California Department of Water Resources. 1964. Quality of Ground Water in California 1961-62, Part 1: Northern and Central California. California Department of Water Resources. Bulletin 66-62.
- California Department of Water Resources. 1966. Precipitation in the Central Valley. Coordinated Statewide Planning Program. California Department of Water Resources, Sacramento District. Office Report.
- California Department of Water Resources. 1975. California's Ground Water. California Department of Water Resources. Bulletin 118.
- California Department of Water Resources. 1975. Progress Report Sacramento And Redding Basins Groundwater Study. California Department of Water Resources, Northern and Central Districts, in cooperation with the U.S. Geological Survey. Bulletin 118.
- California Department of Water Resources. 1976. Progress Report in Ground Water Development Studies, North Sacramento Valley. California Department of Water Resources, Northern District. Memorandum Report.
- California Department of Water Resources. 1980. Ground Water Basins in California. California Department of Water Resources. Bulletin 118-80.
- California Department of Water Resources. 1987. Progress Report Sacramento and Redding Basins Ground Water Study. California Department of Water Resources, Northern and Central Districts, in cooperation with the U.S. Geological Survey.
- California Department of Water Resources. 1993. Sacramento Valley Basin Groundwater Levels -- Butte County. California Department of Water Resources, Northern District. District Report.
- California Department of Water Resources. 1994. Butte and Sutter Basins Water Data Atlas. California Department of Water Resources, Northern District. District Report.
- California Department of Water Resources. 1995. Sacramento Valley Groundwater Quality Investigation. California Department of Water Resources, Northern District.
- California Department of Water Resources. 1998. California Water Plan Update. California Department of Water Resources. Bulletin 160-98, Volumes 1 and 2.
- California Department of Public Works. 1950. Views and Recommendations of State of California on Proposed Report of Chief of Engineers, U.S. Army, on Butte Creek and Cherokee Canal, California. Sacramento: California. Department of Public Works.
- California Division of Water Resources. 1940. Butte Creek Adjudication. Butte Creek and Tributaries Above Western Dam, Butte County, California. Sacramento.
- California Division of Water Resources. 1942. Butte Creek Adjudication. Sacramento: 74 p.
- California Reclamation Board. 1986. Butte Basin Overflow Area Plan of Flood Control: Draft; Project Proposal and Environmental Impact Report. Sacramento: Department of Water Resources Reclamation Board.
- Cherven VB, Edmondson WF. 1992. Structural Geology of the Sacramento Basin: Annual Meeting, Pacific Section AAPG, Sacramento, California, April 27, 1992 - May 2, 1992.
- Dickinson WR, Ingersoll RV, Graham SA. 1979. Paleogene Sediment Dispersal and Paleotectonics in Northern California. Geological Society of America: Bulletin 90:1458-1528.
- Fogelman RP. 1976. Descriptions and Chemical Analysis for Selected Wells in the Central Sacramento Valley, California. USGS. OF-76-472.
- Fogelman RP. 1978. Chemical Quality of Ground Water in the Central Sacramento Valley, California. USGS. Water Resources Investigations 77-133.
- Fogelman RP. 1982. Dissolved-solids Concentrations of Groundwater in the Sacramento Valley, California. USGS. HA-645.

- Fogelman RP. 1983. Ground Water Quality in the Sacramento Valley, California, Water Types and Potential Nitrate and Boron Problem Areas. USGS. HA-651.
- Fogelman RP, Rockwell GL. 1977. Descriptions and Chemical Analysis for Selected Wells in the Eastern Sacramento Valley, California. USGS. OF-77-486.
- Fogleman RP. 1979. Chemical Quality of Ground Water in the Eastern Sacramento Valley, California. USGS.
- Harwood DS, Helley EJ. 1982. Preliminary Structure Contour Map of the Sacramento Valley, California, Showing Major Late Cenozoic Structural Features and Depth to Basement. USGS.
- Harwood DS, Helley EJ. 1987. Late Cenozoic Tectonism of the Sacramento Valley. USGS.
- Harwood DS, Helley EJ, Doukas MP. 1981. Geologic Map of the Chico Monocline and Northeastern Part of the Sacramento Valley, California. USGS.
- Harwood DS, Murchey BL. 1990. Biostratigraphic, Tectonic, and Paleogeographic Ties Between Upper Paleozoic Volcanic and Basinal Rocks in the Northern Sierra Terrane, California. Geological Society of America Special Paper.
- Hawkins FF, Anderson L. 1985. Late Quaternary Tectonics of Part of the Northern Sierra Nevada, California. Geological Society of America.
- Hill KA, Webber JD. 1999. Butte Creek Spring-run Chinook Salmon, *Oncorhynchus tshawytscha*, Juvenile Outmigration and Life History, 1995-1998. Sacramento: State of California Resources Agency Department. of Fish and Game. 46 p.
- Hull LC. 1984. Geochemistry of Groundwater in the Sacramento Valley, California. Central Valley of California RASA Project. USGS. Professional Paper 1401-B.
- Lydon PA. 1969. Geology and Lahars of the Tuscan Formation, Northern California. The Geological Society of America.
- Mankinen EA. 1978. Paleomagnetic Evidence for a Late Cretaceous Deformation of the Great Valley Sequence, Sacramento Valley, California. USGS.
- Mitten HT. 1972. Estimated Ground-water Pumpage in the Northern Part of the Sacramento Valley, California, 1966-69. USGS.
- Mitten HT. 1973. Estimated Ground-water Pumpage in the Northern Part of the Sacramento Valley, California, 1970-71. USGS.
- Page RW. 1974. Base and Thickness of the Post-Eocene Continental Deposits in the Sacramento Valley, California. U.S. Geological Survey in cooperation with California Department of Water Resources. Water Resources Investigations 45-73.
- Page RW. 1986. Geology of the Fresh Groundwater Basin of the Central Valley, California, with Texture Maps and Sections. Regional Aquifer System Analysis. USGS. Professional Paper 1401-C.
- Planert M, Williams JS. 1995. Ground Water Atlas of the United States, Segment 1, California, Nevada. USGS. HA-730-B.
- Poland JF, Evenson RE. 1966. Hydrogeology and Land Subsidence, Great Central Valley, California, Geology of Northern California. California Division of Mines and Geology. 239-247 p.
- Russell RD. 1931. The Tehama Formation of Northern California [Ph.D]: University of California.
- Saucedo GJ, Wagner DL. 1992. Geologic Map of the Chico Quadrangle, California. California Division of Mines and Geology.
- United States Army Corps of Engineers Sacramento District. 1957. Little Chico - Butte Creeks General Design. Sacramento, Calif.: United States Army Corps of Engineers Sacramento District.
- U.S. Geological Survey. 1981. Water Resources Data for California; Volume 4, Northern Central Valley Basins and the Great Basin from Honey Lake Basin to Oregon State Line. USGS.

Williamson AK, Prudic DE, Swain LA. 1985. Groundwater Flow in the Central Valley, California. USGS. OF-85-345.

Williamson AK, Prudic DE, Swain LA. 1989. Groundwater Flow in the Central Valley, California. Regional Aquifer-System Analysis--Central Valley, California. USGS. Professional Paper 1401-D.

## **Errata**

Changes made to the basin description will be noted here.