

Appendix G: Supplemental Water Supply Information

- DWR Groundwater Bulletin 118



CALIFORNIA'S GROUNDWATER

BULLETIN 118 - UPDATE 2003

San Francisco Bay Hydrologic Region




Figure 27 San Francisco Bay Hydrologic Region

Basins and Subbasins of the San Francisco Bay Hydrologic Region

Basin/subbasin	Basin name
2-1	Petaluma Valley
2-2	Napa-Sonoma Valley
2-2.01	Napa Valley
2-2.02	Sonoma Valley
2-2.03	Napa-Sonoma Lowlands
2-3	Suisun-Fairfield Valley
2-4	Pittsburg Plain
2-5	Clayton Valley
2-6	Ygnacio Valley
2-7	San Ramon Valley
2-8	Castro Valley
2-9	Santa Clara Valley
2-9.01	Niles Cone
2-9.02	Santa Clara
2-9.03	San Mateo Plain
2-9.04	East Bay Plain
2-10	Livermore Valley
2-11	Sunol Valley
2-19	Kenwood Valley
2-22	Half Moon Bay Terrace
2-24	San Gregorio Valley
2-26	Pescadero Valley
2-27	Sand Point Area
2-28	Ross Valley
2-29	San Rafael Valley
2-30	Novato Valley
2-31	Arroyo Del Hambre Valley
2-32	Visitacion Valley
2-33	Islais Valley
2-35	Merced Valley
2-36	San Pedro Valley
2-37	South San Francisco
2-38	Lobos
2-39	Marina
2-40	Downtown San Francisco

Description of the Region

The San Francisco Bay HR covers approximately 2.88 million acres (4,500 square miles) and includes all of San Francisco and portions of Marin, Sonoma, Napa, Solano, San Mateo, Santa Clara, Contra Costa, and Alameda counties (Figure 27). The region corresponds to the boundary of RWQCB 2.

Significant geographic features include the Santa Clara, Napa, Sonoma, Petaluma, Suisun-Fairfield, and Livermore valleys; the Marin and San Francisco peninsulas; San Francisco, Suisun, and San Pablo bays; and the Santa Cruz Mountains, Diablo Range, Bolinas Ridge, and Vaca Mountains of the Coast Range. While being the smallest in size of the 10 HRs, the region has the second largest population in the State at about 5.8 million in 1995 (DWR 1998). Major population centers include the cities of San Francisco, San Jose and Oakland.

Groundwater Development

The region has 28 identified groundwater basins. Two of those, the Napa-Sonoma Valley and Santa Clara Valley groundwater basins, are further divided into three and four subbasins, respectively. The groundwater basins underlie approximately 896,000 acres (1,400 square miles) or about 30 percent of the entire HR.

Despite the tremendous urban development in the region, groundwater use accounts for only about 5 percent (68,000 acre-feet) of the region's estimated average water supply for agricultural and urban uses, and accounts for less than one percent of statewide groundwater uses.

In general, the freshwater-bearing aquifers are relatively thin in the smaller basins and moderately thick in the more heavily utilized basins. The more heavily utilized basins in this region include the Santa Clara Valley, Napa-Sonoma Valley, and Petaluma Valley groundwater basins. In these basins, the municipal and irrigation wells have average depths ranging from about 200 to 500 feet. Well yields in these basins range from less than 50 gallons per minute (gpm) to approximately 3,000 gpm. In the smaller basins, most municipal and irrigation wells have average well depths in the 100- to 200-foot range. Well yields in the smaller and less utilized basins are typically less than 500 gpm.

Land subsidence has been a significant problem in the Santa Clara Valley Groundwater Basin in the past. An extensive annual monitoring program has been set up within the basin to evaluate changes in an effort to maintain land subsidence at less than 0.01 feet per year (SCVWD 2001). Additionally, groundwater recharge projects have been implemented in the Santa Clara Valley to ensure that groundwater will continue to be a viable water supply in the future.

Groundwater Quality

In general, groundwater quality throughout most of the region is suitable for most urban and agricultural uses with only local impairments. The primary constituents of concern are high TDS, nitrate, boron, and organic compounds.

The areas of high TDS (and chloride) concentrations are typically found in the region's groundwater basins that are situated close to the San Francisco Bay, such as the northern Santa Clara, southern Sonoma, Petaluma, and Napa valleys. Elevated levels of nitrate have been detected in a large percentage of private wells tested within the Coyote Subbasin and Llagas Subbasin of the Gilroy-Hollister Valley Groundwater Basin (in the Central Coast HR) located to the south of the Santa Clara Valley (SCVWD 2001). The shallow aquifer zone within the Petaluma Valley also shows persistent nitrate contamination. Groundwater with high TDS, iron, and boron levels is present in the Calistoga area of Napa Valley, and elevated boron levels in other parts of Napa Valley make the water unfit for agricultural uses. Releases of fuel hydrocarbons from leaking underground storage tanks and spills/leaks of organic solvents at industrial sites have caused minor to significant groundwater impacts in many basins throughout the region. Methyl tertiary-butyl ether (MTBE) and chlorinated solvent releases to soil and groundwater continue to be problematic. Environmental oversight for many of these sites is performed either by local city and county enforcement agencies, the RWQCB, the Department of Toxic Substances Control, and/or the U.S. Environmental Protection Agency.

Water Quality in Public Supply Wells

From 1994 through 2000, 485 public supply water wells were sampled in 18 of the 33 basins and subbasins in the San Francisco Bay HR. Analyzed samples indicate that 410 wells, or 85 percent, met the state primary MCLs for drinking water standards. Seventy-five wells, or 15 percent, have constituents that exceed one or more MCL. Figure 28 shows the percentages of each contaminant group that exceeded MCLs in the 75 wells.

Table 16 lists the three most frequently occurring contaminants in each contaminant group and the number of wells in the HR that exceeded the MCL for those contaminants.




Figure 28 MCL exceedances in public supply wells in the San Francisco Bay Hydrologic Region

Table 16 Most frequently occurring contaminants by contaminant group in the San Francisco Bay Hydrologic Region

Contaminant group	Contaminant - # of wells	Contaminant - # of wells	Contaminant - # of wells
Inorganics	Iron – 57	Manganese – 57	Fluoride – 7
Radiological	Gross Alpha – 2	Radium 226 – 1	
Nitrates	Nitrate (as NO ₃) – 27	Nitrate + Nitrite – 3	Nitrite (as N) – 1
Pesticides	Di(2-Ethylhexyl)phthalate – 4	Heptachlor – 1	
VOCs/SVOCs	PCE – 4	Dichloromethane – 3	TCE – 2 Vinyl Chloride – 2

TCE = Trichloroethylene
PCE = Tetrachloroethylene
VOC = Volatile Organic Compound
SVOC = Semivolatile Organic Compound

Changes from Bulletin 118-80

Since Bulletin 118-80 was published, RWQCB 2 boundary has been modified. This resulted in several basins being reassigned to RWQCB 1. These are listed in Table 17.

Table 17 Modifications since Bulletin 118-80 of groundwater basins in San Francisco Bay Hydrologic Region

Basin name	New number	Old number
McDowell Valley	1-56	2-12
Knights Valley	1-50	2-13
Potter Valley	1-51	2-14
Ukiah Valley	1-52	2-15
Sanel Valley	1-53	2-16
Alexander Valley	1-54	2-17
Santa Rosa Valley	1-55	2-18
Lower Russian River Valley	1-60	2-20
Bodega Bay Area	1-57	2-21

No additional basins were assigned to the San Francisco Bay HR in this revision. However, the Santa Clara Valley Groundwater Basin (2-9) has been subdivided into four subbasins instead of two, and the Napa-Sonoma Valley Groundwater Basin is now three subbasins instead of two.

There are several deletions of groundwater basins from Bulletin 118-80. The San Francisco Sand Dune Area (2-34) was deleted when the San Francisco groundwater basins were redefined in a USGS report in the early 1990s. The Napa-Sonoma Volcanic Highlands (2-23) is a volcanic aquifer and was not assigned a basin number in this bulletin. This is considered to be a groundwater source area as discussed in Chapter 6. Bulletin 118-80 identified seven groundwater basins that were stated to differ from 118-75: Sonoma County Basin, Napa County Basin, Santa Clara County Basin, San Mateo Basin, Alameda Bay Plain Basin, Niles Cone Basin, and Livermore Basin. They were created primarily by combining several smaller basins and subbasins within individual counties. This report does not consider these seven as basins. There is no change in numbering because the basins were never assigned a basin number.

Table 18 San Francisco Bay Hydrologic Region groundwater data

Basin/Subbasin	Basin Name	Area (acres)	Groundwater Budget Type	Well Yields (gpm)		Active Monitoring			TDS (mg/L)	
				Maximum	Average	Levels	Quality	Title 22	Average	Range
2-1	PETALUMA VALLEY	46,100	C	100	-	16	7	24	347	58-650
2-2	NAPA-SONOMA VALLEY									
2-2.01	NAPA VALLEY	45,900	A	3,000	223	19	10	23	272	150-370
2-2.02	SONOMA VALLEY	44,700	C	1,140	516	18	9	35	321	100-550
2-2.03	NAPA-SONOMA LOWLANDS	40,500	C	300	98	0	6	9	185	50-300
2-3	SUISUN-FAIRFIELD VALLEY	133,600	C	500	200	21	17	35	410	160-740
2-4	PITTSBURG PLAIN	11,600	C	-	-	-	-	9	-	-
2-5	CLAYTON VALLEY	17,800	C	-	-	-	-	48	-	-
2-6	YGNACIO VALLEY	15,500	C	-	-	-	-	-	-	-
2-7	SAN RAMON VALLEY	7,060	C	-	-	-	-	-	-	-
2-8	CASTRO VALLEY	1,820	C	-	-	-	-	-	-	-
2-9	SANTA CLARA VALLEY									
2-9.01	NILES CONE	57,900	A	3,000	2,000	350	120	20	-	-
2-9.02	SANTA CLARA	190,000	C	-	-	-	10	234	408	200-931
2-9.03	SAN MATEO PLAIN	48,100	C	-	-	-	2	14	407	300-480
2-9.04	EAST BAY PLAIN	77,400	A	1,000	UNK	29	16	7	638	364-1,420
2-10	LIVERMORE VALLEY	69,500	A	-	-	-	-	36	-	-
2-11	SUNOL VALLEY	16,600	C	-	-	-	-	2	-	-
2-19	KENWOOD VALLEY	3,170	C	-	-	-	-	13	-	-
2-22	HALF MOON BAY TERRACE	9,150	C	-	-	5	-	9	-	-
2-24	SAN GREGORIO VALLEY	1,070	C	-	-	-	-	-	-	-
2-26	PESCADERO VALLEY	2,900	C	-	-	3	-	4	-	-
2-27	SAND POINT AREA	1,400	C	-	-	-	-	6	-	-
2-28	ROSS VALLEY	1,770	C	-	-	-	-	-	-	-
2-29	SAN RAFAEL VALLEY	880	C	-	-	-	-	-	-	-
2-30	NOVATO VALLEY	20,500	C	-	-	-	-	1	-	-
2-31	ARROYO DEL HAMBRE VALLEY	790	C	-	-	-	-	-	-	-
2-32	VISITACION VALLEY	880	C	-	-	-	-	-	-	-
2-33	ISLAIS VALLEY	1,550	C	-	-	-	-	-	-	-
2-35	MERCED VALLEY	10,400	C	-	-	-	-	10	-	-
2-36	SAN PEDRO VALLEY	880	C	-	-	-	-	-	-	-
2-37	SOUTH SAN FRANCISCO	2,170	C	-	-	-	-	-	-	-
2-38	LOBOS	2,400	A	-	-	-	-	-	-	-
2-39	MARINA	220	A	-	-	-	-	-	-	-
2-40	DOWNTOWN SAN FRANCISCO	7,600	C	-	-	-	-	-	-	-

gpm - gallons per minute

mg/L - milligram per liter

TDS - total dissolved solids

